

Decreased osteoblasts and increased osteoclasts in rats after coal dust exposure

by Nia Kania

Submission date: 08-Mar-2018 12:18PM (UTC+0700)

Submission ID: 927093240

File name: sed_osteoclasts_in_rats_after_coal_dust_exposure_-_NIA_KANIA.pdf (95.46K)

Word count: 2905

Character count: 15195



22

Decreased osteoblasts and increased osteoclasts in rats after coal dust exposure

Izaak Zoelkarnain Akbar*, Nia Kania**, Bambang Setiawan***,
Nurdiana****, and M. Aris Widodo*****

54

ABSTRACT

Bone remodeling is a physiological process of cortical and trabecular bone reconstruction, with initial bone resorption, by osteoclasts and concurrent bone formation by osteoblasts. Oxidative stress due to coal dust exposure not only found in the lungs, but also in the circulation or systemically. The aim of this study was to determine the effect of oxidative stress on coal dust exposure on the number of osteoblasts and osteoclasts in rats. In this experimental study, four groups were evaluated: control; coal dust exposure at 6.25 mg/m³ for 28 days; coal dust exposure at 12.5 mg/m³ for 28 days; coal dust exposure at 25 mg/m³ for 28 days (all exposures were given daily for one hour). Circulatory oxidative stress was measured by malondialdehyde level. Osteoblast and osteoclast numbers were counted by light microscopic examination of distal femoral cross-sections stained with hematoxylin eosin. This study shows that malondialdehyde levels were significantly increased in coal dust exposure groups, in comparison with the control group (p<0.05). There were also significantly decreased numbers of osteoblasts (p<0.05) and significantly increased numbers of osteoclasts (p<0.05) numbers in coal dust exposure groups, as compared with the control group. No correlations were found between malondialdehyde levels (oxidative stress) and respective numbers of osteoblasts and osteoclasts in all coal dust exposure groups (p>0.05). Coal dust exposure increased malondialdehyde level and osteoclast numbers, and decreased osteoblast numbers, but no correlation was found between oxidative stress (caused by coal dust exposure) and osteoblast and osteoclast numbers.

Key words: Coal dust, subchronic, oxidative stress, osteoblast, osteoclast, rat

*Department of Orthopedic Surgery, **Department of Pathology Anatomy, Ulin General Hospital Medical Faculty, Lambung Mangkurat University, Banjarmasin
***Department of Medical Chemistry, Medical Faculty, Lambung Mangkurat University, Banjarbaru
****Department of Pharmacology, Medical Faculty, Brawijaya University, Malang - East Java

Correspondence

dr. Izaak Zoelkarnain Akbar, SpOT(K)
Department of Orthopedic Surgery, Ulin General Hospital Medical Faculty, Lambung Mangkurat University, Jl. A Yani Km.2 No.43 Banjarmasin - South Kalimantan
Email : izaakakbar@yahoo.co.id

Univ Med 2011;30:73-9.

INTRODUCTION

Osteoblasts are bone-forming cells derived from multipotent mesenchymal precursor stem cells, which are also stem cells

for the bone marrow stroma, chondrocytes, muscle cells, and adipocytes, with mean life span of around 3 weeks.^(1,2) Osteoclasts are multinuclear cells of large size (50 to 100 μm in diameter), originating from the

hematopoietic system,^(1,3) with a life span of up to ~2 weeks.⁽⁴⁾ The precursors of osteoclasts are hematopoietic cells of monocyte/macrophage lineage,^(1,2) and the osteoclasts reach the bones through cell migration of progenitors from adjacent connective tissue.⁽¹⁾

Bone remodeling is a physiological process of cortical and trabecular bone reconstruction, with initial bone resorption by osteoclasts and concurrent bone formation by osteoblasts. This process is coordinated throughout life and is the dominant process in adult skeletons. If the resorption and absorption are quantitatively proportional, the remodeling proceeds in a balanced manner.^(5,6) The remodeling process may be disrupted by various toxic substances from the environment, resulting in disease. Exposure to metals, such as cadmium, aluminum, and lead is a risk factor for osteoporosis.⁽⁷⁾

South Kalimantan is the largest coal producer in Indonesia with mines located throughout the area (Banjar, Tanah Laut, Kotabaru, Tanah Bumbu, Hulu Sungai Tengah, Hulu Sungai Utara, Hulu Sungai Selatan, Tapin, and Tabalong). Coal dust is a complex admixture of several minerals, trace metals, and organic substances other than coal particles.⁽⁸⁾ Coal dust particles deposited in the alveolar epithelium are phagocytosed by alveolar macrophages, subsequently increasing release of H₂O₂ and •O₂, triggering oxidative stress.^(9,10) Oxidative stress from coal dust exposure is not only found in the lungs, but also in the circulation or systemically.⁽¹⁰⁾ High levels of oxidative stress causes apoptotic necrosis.⁽¹¹⁾ The numbers of osteoclasts and osteoblasts depend on two processes, i.e. osteoclastogenesis or osteoblastogenesis, and apoptosis.⁽⁴⁾ The study by Parhan et al. demonstrated that lipid oxidation inhibits osteoblast differentiation, thus disturbing bone formation.⁽¹²⁾ *In vitro* and *in vivo* it was also demonstrated that free radicals are involved in osteoclastogenesis and bone resorption.⁽¹³⁾

Therefore the aim of our study was to evaluate the effect of oxidative stress, induced by coal dust exposure, on the numbers of osteoblasts and osteoclasts in rats.

METHODS

Research design

This was a controlled experimental study conducted in the period of October-December 2010.

Experimental animals

The experimental subjects were 2-3 month old male Wistar rats weighing 200-250 grams, subdivided into 4 groups: group 1 was the control group (P1) without exposure to coal dust; group 2 was exposed to coal dust at a dose of 6.25 mg/m³ for 1 hour per day for 28 days (P2); group 3 was exposed to coal dust at a dose of 12.5 mg/m³ for 1 hour per day for 28 days (P3); and group 4 was exposed to coal dust at a dose of 25 mg/m³ for 1 hour per day for 28 days (P4). The sample size was calculated from the formula of Stell et al.⁽¹⁴⁾ to be 6 rats per group. All male Wistar rats were obtained from the Pharmacology Laboratory, Faculty of Medicine, Brawijaya University, Malang.

Preparation of coal dust

Coal dust was prepared by pulverizing coal particles by means of equipment consisting of ball mills, ring mills and Raymond mills at PT. Carsurin Coal Laboratories, Banjarmasin. The resulting coal dust was <75 µm in diameter, and subsequently further screened, using microSieve (BioDesign, USA), producing coal dust of <10 µm diameter as respirable particulate matter.

Exposure to coal dust

Exposure to coal dust was performed by means of coal dust exposure device of 0.5 m³ capacity, designed and provided by the Pharmacology Laboratory, Faculty of Medicine,

Brawijaya University. This device works on the principle of providing an ambient environment containing coal dust as particulate matter, which can be inhaled into the airways of the experimental animals. The air flow through the blower is 1.5-2 liter/minute, which corresponds to the air flow in collieries. The dose of coal dust exposure had been determined by preliminary studies to be 12.5 mg/m³ for a high dose.⁽¹³⁾

Parameter measurements

After undergoing coal dust exposure for 28 days, the experimental animals were sectioned. The rats were anesthetized by means of ether on cotton wool in a plastic container. Sectioning was performed on rats with beating hearts by opening the abdomen, cutting the ribs, and opening the thoracic cavity to find the heart. Blood was obtained from the heart to be used for parameter measurements. The parameters measured in the blood were malondialdehyde (MDA) concentration, performed in the Pharmacology Laboratory, Faculty of Medicine, Brawijaya University, Malang. In addition to the blood samples, the

posterior extremities were also taken for histological sections of the femoral bones with hematoxylin-eosin staining. Subsequently the numbers of osteoblasts and osteoclasts were counted in the Pathology Laboratory, Faculty of Medicine, Brawijaya University, Malang, by means of image analyzer software. For each preparation, 10 high power fields were examined at 400 x magnification.

Ethical clearance

Ethical clearance for work on experimental animals for this study was issued by the Research Ethics Committee, Faculty of Medicine, Lambung Mangkurat University.

Statistical analysis

The collected data were analyzed using ANOVA, and a p value of p<0.05 indicates a significant differences between intervention groups. Significant results of ANOVA were followed up by a post hoc test. In addition correlation tests were performed. All statistical analyses were performed using SPSS software (15.0) for Windows. Minimal significance was assumed at p<0.05.




Figure 1. MDA levels according to coal dust exposure groups in rats. P1=control; P2=coal dust exposure at 6.25 mg/m³ for 28 days; P3=coal dust exposure at 12.5 mg/m³ for 28 days; P4= coal dust exposure at 25 mg/m³ for 28 days




Figure 2. Mean number of osteoblasts based on coal dust exposure groups in rats. P1=control; P2=coal dust exposure at 6.25 mg/m³ for 28 days; P3=coal dust exposure at 12.5 mg/m³ for 28 days; P4= coal dust exposure at 25 mg/m³ for 28 days

RESULTS

According to ANOVA results, subchronic exposure to coal dust at various doses significantly increased malondialdehyde levels in comparison with the control group ($p=0.000$). Using post hoc testing, there were significant differences between intervention groups, except between group P2 and control (P1) ($p=0.192$) (Figure 1).

Using ANOVA, coal dust exposure significantly reduced the mean number of

osteoblasts, as compared to the control group ($p=0.000$). On post hoc testing significant differences were found between intervention groups ($p<0.05$) (Figure 2).

Based on ANOVA, coal dust exposure significantly increased the mean number of osteoclasts, as compared to the control group ($p=0.000$). Using post hoc testing significant differences were found between intervention groups, except between P2 and P3 ($p=0.064$) (Figure 3).




Figure 3. Mean number of osteoclasts based on group in rats. P1=control; P2=coal dust exposure at 6.25 mg/m³ for 28 days; P3=coal dust exposure at 12.5 mg/m³ for 28 days; P4= coal dust exposure at 25 mg/m³ for 28 days




Figure 4. Mechanisms of osteoporosis triggered by coal dust. Oxidation products are formed and accumulate in bone or are transformed into more reactive molecules, thus triggering a) direct effects via mechanisms of 1) inhibition of osteoblast differentiation and increased apoptosis of osteoblasts, thus decreasing bone formation; 2) induction of osteoclasts differentiation, thus increasing bone resorption; b) indirect effects via production of proinflammatory cytokines. Modified after Parhami.⁽¹²⁾

According to the results of the Pearson correlation test, there was no significant correlation between MDA levels and osteoblast and osteoclast numbers in the intervention groups ($p > 0.05$).

DISCUSSION

Inhalation of coal dust leads to formation of reactive oxygen species (ROS) through direct and indirect mechanisms. Direct mechanisms involve bioactive components contained in coal dust, while indirect mechanisms form ROS through respiratory bursts during phagocytosis by macrophages.⁽¹⁰⁾ The content of transitional metals, comprising Fe, Cr, Mn, Co, Ni, Cu, Zn, and silicon, may catalyze Fenton reactions to produce reactive oxygen compounds.⁽¹⁵⁾ During phagocytosis of inhaled particles, superoxide radicals are formed, that are subject to spontaneous dismutation to produce hydrogen peroxide. In the presence of transitional metals, the hydrogen peroxide is converted into hydroxyl radicals.⁽¹⁶⁾

In the present study, increased levels of MDA are the result of oxidative stress, due to coal dust exposure. Formation of MDA is by lipid peroxidation, which is significantly different between intervention groups. Increased lipid peroxidation is consistent with the study by Armutcu et al.⁽¹⁰⁾ In addition, according to the hypothesis of Parhami,⁽¹²⁾ lipid peroxidation may affect the numbers of osteoblasts and osteoclasts in the aged condition (Figure 4). In our study, the number of osteoblasts decreased concurrently with increased dose of coal dust exposure, although no significant difference was found between P2 and P3. In addition, the number of osteoclasts increased concurrently with increased doses of coal dust exposure, although no significant difference was found between P2 and P3. This indicates an imbalance between bone formation and resorption.

Interestingly, there was no correlation between MDA levels and the number of osteoblasts, or between MDA levels and the numbers of osteoclasts, which was not according to Parhami's hypothesis.⁽¹²⁾ This is presumably due to the presence of other

molecules capable of affecting the numbers of osteoblasts and osteoclasts. MDA molecules are active molecules that may undergo transformation into other molecules, e.g. via enzymatic glycosylation. The available aldehyde groups may react with amino groups of proteins, giving rise to advanced glycation end products (AGEs), which subsequently may affect osteoblasts and osteoclasts.⁽¹⁷⁻¹⁹⁾ Another mechanism of coal dust is through triggering of increases in nitrous oxide levels, which may trigger apoptosis of osteoblasts.^(10,20)

One limitation of this study is the failure to explore the various parameters that are increased as a result of coal dust exposure, in order to demonstrate the markers that are the cause of the reduction in the numbers of osteoblasts and the increased numbers of osteoclasts.

CONCLUSION

Exposure to coal dust increases MDA levels and the numbers of osteoclasts, while reducing the numbers of osteoblasts, although there was no significant correlation between oxidative stress due to coal dust exposure and the numbers of osteoblasts and osteoclasts.

ACKNOWLEDGEMENT

We thank PT. Carsurin Coal Laboratories for the supply of coal dust of <75 µm diameter.

REFERENCES

1. Manolagas SC. Birth and death of bone cells: basic regulatory mechanisms and implications for the pathogenesis and treatment of osteoporosis. *Proc Natl Acad Sci U S A* 2000;21:115-37.
2. Okamoto M, Murai J, Yoshikawa H, Tsumaki N. Bone morphogenetic proteins in bone stimulate osteoclasts and osteoblasts during bone development. *J Bone Min Res* 2006;21:1022-33.
3. Saltel F, Chabadel A, Zhao Y, Lafage-Proust M, Clezardin P, Jurdic P, et al. Transmigration: a new property of mature multinucleated osteoclasts. *J Bone Min Res* 2006;21:1913-23.
4. Wu X, Pan G, McKenna MA, Zayzafoon M, Xiong W, McDonald JM. RANKL regulates fas expression and fas-mediated apoptosis in osteoclasts. *J Bone Min Res* 2005;20:107-16.
5. Lemer UH. Bone remodeling in postmenopausal osteoporosis. *J Dent Res* 2006;85:584.
6. Epstein S. The roles of bone mineral density, bone turnover, and other properties in reducing fracture risk during antiresorptive therapy. *Mayo Clin Proc* 2005;80:379-88.
7. Alfvén T, Elinder C, Carlsson MD, Grubb A, Hellström L, Persson B, et al. Low-level cadmium exposure and osteoporosis. *J Bone Min Res* 2000;15:1579-86.
8. Dalal NS, Newman J, Pack D, Leonard S, Valyathan V. Hydroxyl radical generation by coal mine dust: possible implication to coal workers' pneumoconiosis. *Free Radic Biol Med* 1995;18:220.
9. Huang X, Finkelman RB. Understanding the chemical properties of macerals and minerals in coal and its potential application for occupational lung disease prevention. *J Toxicol Environ Health Part B* 2008;1:45-67.
10. Armutcu F, Gun BD, Altin R, Gurel A. Examination of lung toxicity, oxidant/antioxidant status and effect of erdosteine in rats kept in coal mine ambience. *Environ Toxicol Pharmacol* 2007;24:106-13.
11. Glorie G, Legrand-Poels S, Piette J. NF-κB activation by reactive oxygen species: fifteen years later. *Biochem Pharmacol* 2006;72:1493-20.
12. Parhami F. Possible role of oxidized lipids in osteoporosis: could hyperlipidemia be a risk factor? *Prostaglandins Leukot Essent Fatty Acids* 2003;68:373-8.
13. Ghanem MM, Porter D, Batteli LA, Valyathan V, Kashon ML, Ma JY, et al. Respirable coal dust particle modify cytochrome P4501A1 expression in rat alveolar cells. *Am J Resp Cell Mol Biol* 2004;31:171-87.
14. Steel RGD, Torrie JH, Dickey DA. Principles and procedures of statistics, a biometrical approach. 3rd ed. Singapore: McGraw Hill Book Company; 1997.
15. Nadif R, Mintz M, Jedlicka A, Bertrand J, Kleiberger SR, Kauffmann F. Association of CAT polymorphisms with catalase activity and exposure to environmental oxidative stimuli. *Free Radic Res* 2005;39:1345-50.
16. Altin R, Kart L, Tekin I, Armutcu F, Tor M, Ornek T. The presence of promatrix metalloproteinase-3 and its relation with different categories of coal worker's pneumoconiosis. *Mediators Inflamm* 2004;13:105-9.

17. ³ Hein G, Wiegand R, Lehmann G, Stein G, Franke S. Advanced glycation end-products pentosidine and N^ε-carboxymethyllysine are elevated in serum of patients with osteoporosis. *Rheumatol* 2003; 42:1242-6.
18. Franke S, Siggelkow H, Wolf G, Hein G. Advanced glycation endproducts influence the mRNA expression of RAGE, RANKL and various osteoblastic genes in human osteoblasts. *Arch Physiol Biochem* 2007;113:154-61.
19. ⁵ Alikhani M, Alikhani Z, Boyd C, MacLellan CM, Raptis M, Liu R, et al. Advanced glycation end products stimulate osteoblast apoptosis via the MAP kinase and cytosolic apoptotic pathways. *Journal of Cellular Biochemistry* 2007;40:345-53.
20. ¹⁹ Chen RM, Chen TL, Chiu WT, Chang CC. Molecular mechanism of nitric oxide-induced osteoblast apoptosis. *J Orthop Res* 2005;23:462-8.

Decreased osteoblasts and increased osteoclasts in rats after coal dust exposure

ORIGINALITY REPORT

44%

SIMILARITY INDEX

28%

INTERNET SOURCES

41%

PUBLICATIONS

22%

STUDENT PAPERS

PRIMARY SOURCES

- 1** "IOF Regionals—1st Asia-Pacific Osteoporosis Meeting : Poster Presentation Abstracts", Osteoporosis International, 12/2010
Publication **5%**
- 2** oamsjournal.com
Internet Source **3%**
- 3** Akbar, Izaak, Nur Permatasari, Djoko Soeatmadji, Mohammad Hidayat, and Handono Kalim. "Effect of methylglyoxal on osteoprotegerin expression in the pre-osteoblast MC3T3E1 cell line: involvement of oxidant pathways", Journal of Experimental and Integrative Medicine, 2013.
Publication **2%**
- 4** "Poster Presentations", FEBS Journal, 6/2006
Publication **2%**
- 5** diabeticfootandankle.net
Internet Source **1%**

Mohamed M. Ghanem, Lori A. Battelli, Robert

6

R. Mercer, James F. Scabilloni, Michael L. Kashon, Jane Y.C. Ma, Joginder Nath, Ann F. Hubbs. "Apoptosis and Bax Expression are Increased by Coal Dust in the Polycyclic Aromatic Hydrocarbon-Exposed Lung", *Environmental Health Perspectives*, 2006

Publication

1%

7

jeim.org

Internet Source

1%

8

www.ajcn.org

Internet Source

1%

9

www.jdmdonline.com

Internet Source

1%

10

tipfakultesi.bezmialem.edu.tr

Internet Source

1%

11

osteoporoza.meditorial.cz

Internet Source

1%

12

Gruber, R.. "BMP-6-induced osteogenic differentiation of mesenchymal cell lines is not modulated by sex steroids and resveratrol", *Cytokine*, 20030821/0907

Publication

1%

13

www.uasom.uab.edu

Internet Source

1%

14

Delaisse, Jean-Marie. "The reversal phase of

1%

the bone-remodeling cycle: cellular prerequisites for coupling resorption and formation", BoneKEy Reports, 2014.

Publication

15

Systems Biology of Free Radicals and Antioxidants, 2014.

Publication

1%

16

etheses.whiterose.ac.uk

Internet Source

1%

17

www.ftrdergisi.com

Internet Source

1%

18

Kania, Nia, Bambang Setiawan, Edi Widjadjanto, Nurdiana Nurdiana, M. Aris Widodo, and H.M.S. Chandra Kusuma. "Subchronic inhalation of coal dust particulate matter 10 induces bronchoalveolar hyperplasia and decreases MUC5AC expression in male Wistar rats", Experimental and Toxicologic Pathology, 2014.

Publication

1%

19

www.cellbiolint.org

Internet Source

1%

20

Nicole E Jensky. "The association of bone density and calcified atherosclerosis is stronger in women without dyslipidemia: The multi-ethnic study of atherosclerosis", Journal of

1%

Bone and Mineral Research, 11/2011

Publication

| | | |
|----|--|----|
| 21 | www.psy.uq.edu.au Internet Source | 1% |
| 22 | www.library.gunadarma.ac.id Internet Source | 1% |
| 23 | www.bumc.bu.edu Internet Source | 1% |
| 24 | baranyilaszlozsolt.com Internet Source | 1% |
| 25 | soar.wichita.edu Internet Source | 1% |
| 26 | "European Congress on Osteoporosis & Osteoarthritis (IOF-ECCEO12) : Poster Presentation Abstracts", Osteoporosis International, 03/2012 Publication | 1% |
| 27 | Submitted to Universitas Brawijaya Student Paper | 1% |
| 28 | ptjournal.org Internet Source | 1% |
| 29 | help.acdsystems.com Internet Source | 1% |
| 30 | Setiawan, Bambang, Asnawati Darsuni, Fauzan | 1% |

Muttaqien, Dwi Adiputro, Nia Kania, Dian Nugrahenny, and M Widodo. "The effects of combined particulate matter 10 coal dust exposure and high-cholesterol diet on lipid profiles, endothelial damage, and hematopoietic stem cells in rats", *Journal of Experimental and Integrative Medicine*, 2013.

Publication

31

Submitted to Miami University of Ohio

Student Paper

1%

32

nurmelatisseptiana.blogspot.com

Internet Source

1%

33

Armutcu, F.. "Examination of lung toxicity, oxidant/antioxidant status and effect of erdosteine in rats kept in coal mine ambience", *Environmental Toxicology and Pharmacology*, 200709

Publication

1%

34

Kania, Nia, Bambang Setiawan, Edi Widjajanto, Nurdiana Nurdiana, M Widodo, and HMS Kusuma. "Peroxidative index as novel marker of hydrogen peroxide involvement in lipid peroxidation from coal dust exposure", *Oxidants and Antioxidants in Medical Science*, 2012.

Publication

1%

Jorge B Cannata-Andía. "Estrogens and bone

35

disease in chronic kidney disease: role of FGF23 :", Current Opinion in Nephrology & Hypertension, 07/2010

Publication

<1%

36

Sakae Tanaka. "Role of RANKL in physiological and pathological bone resorption and therapeutics targeting the RANKL-RANK signaling system", Immunological Reviews, 12/2005

Publication

<1%

37

Submitted to Shenandoah University

Student Paper

<1%

38

Setiawan, Bambang, Asnawati Darsuni, Fauzan Muttaqien, and M Widodo. "Cholesterol Lowering Effect of Subchronic Inhalation Particulate Matter 10 Coal Dust on Rats", Medicine Science | International Medical Journal, 2013.

Publication

<1%

39

Itoh, S.. "A critical role for interleukin-6 family-mediated Stat3 activation in osteoblast differentiation and bone formation", Bone, 200609

Publication

<1%

40

Noor, Zairin, and Bambang Setiawan. "Combined inhalation of cigarette smoke and coal dust particulate matter 10 increase bone

<1%

iron level in rats", Journal of Experimental and Integrative Medicine, 2012.

Publication

41

www.mkb-online.org

Internet Source

<1%

42

Ulrich Valcourt. "BMP Signaling in Osteogenesis, Bone Remodeling and Repair", European Journal of Trauma, 10/2005

Publication

<1%

43

Xi Huang. "Understanding the Chemical Properties of Macerals and Minerals in Coal and its Potential Application for Occupational Lung Disease Prevention", Journal of Toxicology and Environmental Health Part B, 01/2008

Publication

<1%

44

McCormick, R. Keith. "Osteoporosis: integrating biomarkers and other diagnostic correlates into the management of bone fra", Alternative Medicine Review, June 2007 Issue

Publication

<1%

45

www.stat.osu.edu

Internet Source

<1%

46

Noor, Zairin, and Bambang Setiawan. "Subchronic inhalation of coal dust particulate matter 10 changes bone mesostructure, mineral element levels and turnover markers in

<1%

rats", Journal of Experimental and Integrative
Medicine, 2013.

Publication

47

www.sld.cu

Internet Source

<1%

48

content.karger.com

Internet Source

<1%

49

Yuqiong Li, Jianhua Chen, Ye Chen, Cuihua
Zhao, Yibing Zhang, Baolin Ke. "Interactions of
Oxygen and Water Molecules with Pyrite
Surface: A New Insight", Langmuir, 2018

Publication

<1%

50

Oxidative Stress and Free Radical Damage in
Neurology, 2011.

Publication

<1%

51

S. Ávila Júnior. "Occupational airborne
contamination in south Brazil: 1. Oxidative
stress detected in the blood of coal miners",
Ecotoxicology, 07/18/2009

Publication

<1%

52

ajrccm.atsjournals.org

Internet Source

<1%

53

Principles of Osteoimmunology, 2012.

Publication

<1%

54

Paul, Sangita, John C. Lee, and Lee-Chuan C.

<1%

Yeh. "A comparative study on BMP-induced osteoclastogenesis and osteoblastogenesis in primary cultures of adult rat bone marrow cells", Growth Factors, 2009.

Publication

55

Roux, S.. "Death receptors, Fas and TRAIL receptors, are involved in human osteoclast apoptosis", Biochemical and Biophysical Research Communications, 20050722

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off